

## HÖLDER TYPE INEQUALITIES ON HILBERT C\*-MODULES AND ITS REVERSES

YUKI SEO<sup>1</sup>

*This paper is dedicated to Professor Tsuyoshi Ando*

ABSTRACT. In this paper, we show Hilbert C\*-module versions of Hölder-McCarthy inequality and its complementary inequality. As an application, we obtain Hölder type inequalities and its reverses on a Hilbert C\*-module.

### 1. INTRODUCTION

The Hölder inequality is one of the most important inequalities in functional analysis. If  $a = (a_1, \dots, a_n)$  and  $b = (b_1, \dots, b_n)$  are  $n$ -tuples of nonnegative numbers, and  $1/p + 1/q = 1$ , then

$$\sum_{i=1}^n a_i b_i \leq \left( \sum_{i=1}^n a_i^p \right)^{1/p} \left( \sum_{i=1}^n b_i^q \right)^{1/q} \quad \text{for all } p > 1$$

and

$$\sum_{i=1}^n a_i b_i \geq \left( \sum_{i=1}^n a_i^p \right)^{1/p} \left( \sum_{i=1}^n b_i^q \right)^{1/q} \quad \text{for all } p < 0 \text{ or } 0 < p < 1.$$

Non-commutative versions of the Hölder inequality and its reverses have been studied by many authors. T. Ando [1] showed the Hadamard product version of a Hölder type. T. Ando and F. Hiai [2] discussed the norm Hölder inequality and the matrix Hölder inequality. B. Mond and O. Shisha [15], M. Fujii, S. Izumino, R. Nakamoto and Y. Seo [7], and S. Izumino and M. Tominaga [11] considered the vector state version of a Hölder type and its reverses. J.-C. Bourin, E.-Y. Lee, M. Fujii and Y. Seo [3] showed the geometric operator mean version, and so on.

---

2010 *Mathematics Subject Classification.* Primary 46L08; Secondary 47A63 and 47A64.

*Key words and phrases.* Hölder-McCarthy inequality, Hölder inequality, Hilbert C\*-modules, geometric operator mean.

In this paper, as a generalization of the vector state version due to [7], we show Hilbert C\*-module versions of Hölder-McCarthy inequality and its complementary inequality. As an application, we obtain Hölder type inequalities and its reverses on a Hilbert C\*-module.

## 2. PRELIMINARY

Let  $\mathcal{B}(H)$  be the C\*-algebra of all bounded linear operators on a Hilbert space  $H$ , and  $\mathcal{A}$  be a unital C\*-algebra of  $\mathcal{B}(H)$  with the unit element  $e$ . For  $a \in \mathcal{A}$ , we denote the *absolute value* of  $a$  by  $|a| = (a^*a)^{\frac{1}{2}}$ . For positive elements  $a, b \in \mathcal{A}$  and  $t \in [0, 1]$ , the *t-geometric mean* of  $a$  and  $b$  in the sense of Kubo-Ando theory [12] is defined by

$$a \sharp_t b = a^{\frac{1}{2}} \left( a^{-\frac{1}{2}} b a^{-\frac{1}{2}} \right)^t a^{\frac{1}{2}}$$

for  $a > 0$ , i.e.,  $a$  is invertible. In the case of non-invertible, since  $a \sharp_t b$  satisfies the upper semicontinuity, we define  $a \sharp_t b = \lim_{\varepsilon \rightarrow +0} (a + \varepsilon e) \sharp_t (b + \varepsilon e)$  in the strong operator topology. Hence  $a \sharp_t b \in \mathcal{A}''$  in general, where  $\mathcal{A}''$  is the bi-commutant of  $\mathcal{A}$ . In the case of  $t = 1/2$ , we denote  $a \sharp_{1/2} b$  by  $a \sharp b$  simply. The operator geometric mean has the symmetric property:  $a \sharp_t b = b \sharp_{1-t} a$ , and  $a \sharp_t b = a^{1-t} b^t$  for commuting  $a$  and  $b$ .

A complex linear space  $\mathcal{X}$  is said to be an *inner product  $\mathcal{A}$ -module* (or a pre-Hilbert  $\mathcal{A}$ -module) if  $\mathcal{X}$  is a right  $\mathcal{A}$ -module together with a C\*-valued map  $\langle x, y \rangle \mapsto \langle x, y \rangle : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{A}$  such that

- (i)  $\langle x, \alpha y + \beta z \rangle = \alpha \langle x, y \rangle + \beta \langle x, z \rangle \quad (x, y, z \in \mathcal{X}, \alpha, \beta \in \mathbb{C})$ ,
- (ii)  $\langle x, ya \rangle = \langle x, y \rangle a \quad (x, y \in \mathcal{X}, a \in \mathcal{A})$ ,
- (iii)  $\langle y, x \rangle = \langle x, y \rangle^* \quad (x, y \in \mathcal{X})$ ,
- (iv)  $\langle x, x \rangle \geq 0 \quad (x \in \mathcal{X})$  and if  $\langle x, x \rangle = 0$ , then  $x = 0$ .

The linear structures of  $\mathcal{A}$  and  $\mathcal{X}$  are assumed to be compatible. If  $\mathcal{X}$  satisfies all conditions for an inner-product  $\mathcal{A}$ -module except for the second part of (iv), then we call  $\mathcal{X}$  a *semi-inner product  $\mathcal{A}$ -module*.

Let  $\mathcal{X}$  be an inner product  $\mathcal{A}$ -module over a unital C\*-algebra  $\mathcal{A}$ . We define the norm of  $\mathcal{X}$  by  $\|x\| := \sqrt{\|\langle x, x \rangle\|}$  for  $x \in \mathcal{X}$ , where the latter norm denotes the C\*-norm of  $\mathcal{A}$ . If  $\mathcal{X}$  is complete with respect to this norm, then  $\mathcal{X}$  is called a *Hilbert  $\mathcal{A}$ -module*. An element  $x$  of the Hilbert  $\mathcal{A}$ -module is called *nonsingular* if the element  $\langle x, x \rangle \in \mathcal{A}$  is invertible. For more details on Hilbert C\*-modules, see [13, 14].

In [6], from a viewpoint of operator geometric mean, we showed the following new Cauchy-Schwarz inequality:

**Theorem 2.1** (Cauchy-Schwarz inequality). *Let  $\mathcal{X}$  be a semi-inner product  $\mathcal{A}$ -module over a unital C\*-algebra  $\mathcal{A}$ . If  $x, y \in \mathcal{X}$  such that the inner product  $\langle x, y \rangle$  has a polar decomposition  $\langle x, y \rangle = u|\langle x, y \rangle|$  with a partial isometry  $u \in \mathcal{A}$ , then*

$$|\langle x, y \rangle| \leq u^* \langle x, x \rangle u \sharp \langle y, y \rangle. \quad (2.1)$$

*Under the assumption that  $\mathcal{X}$  is an inner product  $\mathcal{A}$ -module and  $y$  is nonsingular, the equality in (2.1) holds if and only if  $xu = yb$  for some  $b \in \mathcal{A}$ .*

Next we review the basic concepts of adjointable operators on a Hilbert C\*-module. Let  $\mathcal{X}$  be a Hilbert C\*-module over a unital C\*-algebra  $\mathcal{A}$ . Let  $End_{\mathcal{A}}(\mathcal{X})$  denote the set of all bounded  $\mathbb{C}$ -linear  $\mathcal{A}$ -homomorphism from  $\mathcal{X}$  to  $\mathcal{X}$ . Let  $T \in End_{\mathcal{A}}(\mathcal{X})$ . We say that  $T$  is *adjointable* if there exists a  $T^* \in End_{\mathcal{A}}(\mathcal{X})$  such that  $\langle Tx, y \rangle = \langle x, T^*y \rangle$  for all  $x, y \in \mathcal{X}$ . Let  $\mathcal{L}(\mathcal{X})$  denote the set of all adjointable operators from  $\mathcal{X}$  to  $\mathcal{X}$ . Moreover, we define its norm by

$$\|T\| = \sup\{\|\langle Tx, Tx \rangle\|^{\frac{1}{2}} : \|x\| \leq 1\}.$$

Then  $\mathcal{L}(\mathcal{X})$  is a C\*-algebra. The symbol  $I$  stands for the identity operator in  $\mathcal{L}(\mathcal{X})$ . The following lemma due to Pashke [16] is very important:

**Lemma 2.2.** *Let  $\mathcal{X}$  be a Hilbert C\*-module and let  $T$  be a bounded  $\mathcal{A}$ -linear operator on  $\mathcal{X}$ . The following conditions are equivalent:*

- (1)  $T$  is a positive element of  $\mathcal{L}(\mathcal{X})$ ;
- (2)  $\langle x, Tx \rangle \geq 0$  for all  $x$  in  $\mathcal{X}$ .

In [8], we showed the following generalized Cauchy–Schwarz inequality on a Hilbert C\*-module by virtue of (2.1) and Lemma 2.2:

**Theorem 2.3** (generalized Cauchy-Schwarz inequality). *Let  $T$  be a positive operator in  $\mathcal{L}(\mathcal{X})$ . If  $x, y \in \mathcal{X}$  such that  $\langle x, Ty \rangle$  has a polar decomposition  $\langle x, Ty \rangle = u|\langle x, Ty \rangle|$  with a partial isometry  $u \in \mathcal{A}$ , then*

$$|\langle x, Ty \rangle| \leq u^* \langle x, Tx \rangle u \sharp \langle y, Ty \rangle. \quad (2.2)$$

*Under the assumption that  $\langle y, Ty \rangle$  is invertible, the equality in (2.2) holds if and only if  $T^{\frac{1}{2}}(xu) = T^{\frac{1}{2}}(yb)$  for some  $b \in \mathcal{A}$ .*

### 3. HÖLDER-MCCARTHY INEQUALITY

In this section, we show two Hilbert C\*-module versions of Hölder-McCarthy inequality and its complementary inequality. For convenience, we use the notation  $\natural_t$  for the binary operation

$$a \natural_t b = a^{\frac{1}{2}} \left( a^{-\frac{1}{2}} b a^{-\frac{1}{2}} \right)^t a^{\frac{1}{2}} \quad \text{for } t \notin [0, 1],$$

whose formula is the same as  $\sharp_t$ .

**Theorem 3.1.** *Let  $T$  be a positive operator in  $\mathcal{L}(\mathcal{X})$  and  $x$  a nonsingular element of  $\mathcal{X}$ .*

- (1) *If  $p \geq 1$ , then  $\langle x, Tx \rangle \leq \langle x, x \rangle \sharp_{1/p} \langle x, T^p x \rangle$ .*
- (2) *If  $p \leq -1$  or  $1/2 \leq p \leq 1$ , then  $\langle x, x \rangle \natural_{1/p} \langle x, T^p x \rangle \leq \langle x, Tx \rangle$ .*

*Proof.* For a nonsingular element  $x$  of  $\mathcal{X}$ , Put

$$\Phi_x(X) = \langle x \langle x, x \rangle^{-\frac{1}{2}}, X x \langle x, x \rangle^{-\frac{1}{2}} \rangle \quad \text{for } X \in \mathcal{L}(\mathcal{X}).$$

Then  $\Phi_x$  is a unital positive linear map from  $\mathcal{L}(\mathcal{X})$  to  $\mathcal{A}$ .

Suppose that  $p \geq 1$ . Since  $t^{1/p}$  is operator concave, it follows from [4, 5] that  $\Phi_x(T^{1/p}) \leq \Phi_x(T)^{1/p}$  and this implies

$$\langle x, x \rangle^{-\frac{1}{2}} \langle x, T^{1/p} x \rangle \langle x, x \rangle^{-\frac{1}{2}} \leq \left( \langle x, x \rangle^{-\frac{1}{2}} \langle x, Tx \rangle \langle x, x \rangle^{-\frac{1}{2}} \right)^{1/p}$$

and

$$\begin{aligned} \langle x, T^{1/p} x \rangle &\leq \langle x, x \rangle^{\frac{1}{2}} \left( \langle x, x \rangle^{-\frac{1}{2}} \langle x, Tx \rangle \langle x, x \rangle^{-\frac{1}{2}} \right)^{1/p} \langle x, x \rangle^{\frac{1}{2}} \\ &= \langle x, x \rangle \sharp_{1/p} \langle x, Tx \rangle. \end{aligned} \quad (3.1)$$

Replacing  $T$  by  $T^p$  in (3.1), we have (1).

Suppose that  $p \leq -1$  or  $1/2 \leq p \leq 1$ . Since  $-1 \leq 1/p < 0$  or  $1 \leq 1/p \leq 2$ , we have  $\Phi_x(T)^{\frac{1}{p}} \leq \Phi_x(T^{\frac{1}{p}})$  by the operator convexity of  $t^{1/p}$  and this implies

$$\left( \langle x, x \rangle^{-\frac{1}{2}} \langle x, Tx \rangle \langle x, x \rangle^{-\frac{1}{2}} \right)^{\frac{1}{p}} \leq \langle x, x \rangle^{-\frac{1}{2}} \langle x, T^{\frac{1}{p}} x \rangle \langle x, x \rangle^{-\frac{1}{2}}.$$

Hence it follows that

$$\langle x, x \rangle \sharp_{1/p} \langle x, Tx \rangle \leq \langle x, T^{\frac{1}{p}} x \rangle \quad (3.2)$$

and replacing  $T$  by  $T^p$  in (3.2) we have (2).  $\square$

*Remark 3.2.* The inequality (2) of Theorem 3.1 does not hold for  $0 < p < 1/2$  in general. In fact, we give a simple counterexample to the case of  $p = 1/3$  as follows: Put

$$\mathcal{X} = M_4(\mathbb{C}) = M_2(M_2(\mathbb{C})) \quad \text{and} \quad \mathcal{A} = M_2(\mathbb{C}) \oplus M_2(\mathbb{C})$$

and

$$\Phi\left(\begin{pmatrix} X & Y \\ Z & W \end{pmatrix}\right) = \begin{pmatrix} X & 0 \\ 0 & W \end{pmatrix}$$

for  $X, Y, Z, W \in M_2(\mathbb{C})$ . Then  $\mathcal{X}$  is a Hilbert  $\mathcal{A}$ -module with an inner product  $\langle x, y \rangle = \Phi(x^*y)$  for  $x, y \in \mathcal{X}$ . Let

$$z = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix} \quad \text{and} \quad x = \begin{pmatrix} 1/2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

If  $T = T_z$  is defined by  $T_z y = zy$  for all  $y \in \mathcal{X}$ , then  $T$  is a positive operator in  $\mathcal{L}(\mathcal{X})$  and

$$\left( \langle x, x \rangle^{-1/2} \langle x, Tx \rangle \langle x, x \rangle^{-1/2} \right)^3 = \begin{pmatrix} 13 & 8 \\ 8 & 5 \end{pmatrix} \oplus \begin{pmatrix} 4 & 4 \\ 4 & 4 \end{pmatrix}$$

and

$$\langle x, x \rangle^{-1/2} \langle x, T^3 x \rangle \langle x, x \rangle^{-1/2} = \begin{pmatrix} 29 & 22 \\ 22 & 17 \end{pmatrix} \oplus \begin{pmatrix} 17 & 17 \\ 17 & 17 \end{pmatrix},$$

so that

$$\begin{aligned} &\langle x, x \rangle^{-1/2} \langle x, T^3 x \rangle \langle x, x \rangle^{-1/2} - \left( \langle x, x \rangle^{-1/2} \langle x, Tx \rangle \langle x, x \rangle^{-1/2} \right)^3 \\ &= \begin{pmatrix} 16 & 14 \\ 14 & 12 \end{pmatrix} \oplus \begin{pmatrix} 13 & 13 \\ 13 & 13 \end{pmatrix} \not\geq 0 \oplus 0. \end{aligned}$$

Next, we show a complementary part of Theorem 3.1. For this, we need the *generalized Kantorovich constant*  $K(\alpha, \beta, p)$  for  $0 < \alpha < \beta$ , which is defined by

$$K(\alpha, \beta, p) = \frac{\alpha\beta^p - \beta\alpha^p}{(p-1)(\beta-\alpha)} \left( \frac{p-1}{p} \frac{\beta^p - \alpha^p}{\alpha\beta^p - \beta\alpha^p} \right)^p \quad (3.3)$$

for any real number  $p \in \mathbb{R}$ , see also [10, Definition 2.2]. The constant  $K(\alpha, \beta, p)$  satisfies  $0 < K(\alpha, \beta, p) \leq 1$  for  $0 \leq p \leq 1$  and  $K(\alpha, \beta, p) \geq 1$  for  $p \notin [0, 1]$ . For more details on the generalized Kantorovich constant, see [10, Chapter 2.7].

**Theorem 3.3.** *Let  $T$  be a positive invertible operator in  $\mathcal{L}(\mathcal{X})$  such that  $\alpha I \leq T \leq \beta I$  for some scalars  $0 < \alpha < \beta$ , and  $x$  a nonsingular element of  $\mathcal{X}$ .*

(1) *If  $p \geq 1$ , then*

$$\langle x, x \rangle \sharp_{1/p} \langle x, T^p x \rangle \leq K(\alpha, \beta, p)^{1/p} \langle x, Tx \rangle.$$

(2) *If  $p \leq -1$  or  $1/2 \leq p \leq 1$ , then*

$$\langle x, Tx \rangle \leq K(\alpha^p, \beta^p, 1/p) \langle x, x \rangle \natural_{1/p} \langle x, T^p x \rangle,$$

where the generalized Kantorovich constant  $K(\alpha, \beta, p)$  is defined by (3.3).

*Proof.* For a nonsingular element  $x$  of  $\mathcal{X}$ , put  $\Phi_x(X) = \langle x \langle x, x \rangle^{-\frac{1}{2}}, Xx \langle x, x \rangle^{-\frac{1}{2}} \rangle$  for  $X \in \mathcal{L}(\mathcal{X})$ . Then  $\Phi_x : \mathcal{L}(\mathcal{X}) \mapsto \mathcal{A}$  is a unital positive linear map.

Suppose that  $p \geq 1$ . It follows from [10, Lemma 4.3] that

$$\Phi_x(T^p) \leq K(\alpha, \beta, p) \Phi_x(T)^p.$$

This implies

$$\langle x, x \rangle \sharp_{1/p} \langle x, T^p x \rangle \leq K(\alpha, \beta, p)^{1/p} \langle x, Tx \rangle$$

and we have (1).

In the case of  $p \leq -1$  or  $1/2 \leq p \leq 1$ , since  $-1 \leq 1/p < 0$  or  $1 \leq 1/p \leq 2$ , it follows that  $\Phi_x(T^{1/p}) \leq K(\alpha, \beta, 1/p) \Phi_x(T)^{1/p}$ . Similarly we have the desired inequality (2).  $\square$

Next, we discuss Hölder-McCarthy type inequalities on a Hilbert C\*-module outside intervals of Theorem 3.1.

**Corollary 3.4.** *Let  $T$  be a positive invertible operator in  $\mathcal{L}(\mathcal{X})$  such that  $\alpha I \leq T \leq \beta I$  for some scalars  $0 < \alpha < \beta$ , and  $x$  a nonsingular element of  $\mathcal{X}$ . If  $-1 < p < 0$  or  $0 < p < 1/2$ , then*

$$K(\alpha^p, \beta^p, 1/p)^{-1} \langle x, Tx \rangle \leq \langle x, x \rangle \natural_{1/p} \langle x, T^p x \rangle \leq K(\alpha^p, \beta^p, 1/p) \langle x, Tx \rangle,$$

where the generalized Kantorovich constant  $K(\alpha, \beta, p)$  is defined by (3.3).

*Proof.* For a unital positive linear map  $\Phi_x$  from  $\mathcal{L}(\mathcal{X})$  to  $\mathcal{A}$ , it follows from [10, Lemma 4.3] that for  $-1 < p < 0$  or  $0 < p < 1/2$

$$K(\alpha, \beta, 1/p)^{-1} \Phi_x(T)^{1/p} \leq \Phi_x(T^{1/p}) \leq K(\alpha, \beta, 1/p) \Phi_x(T)^{1/p}.$$

Hence we have this theorem as in the proof of Theorem 3.3.  $\square$

Similarly we have the following Hölder-McCarthy type inequality on a Hilbert C\*-module and its complementary inequality as follows:

**Theorem 3.5.** *Let  $T$  be a positive invertible operator in  $\mathcal{L}(\mathcal{X})$  such that  $\alpha I \leq T \leq \beta I$  for some scalars  $0 < \alpha < \beta$ . Then for  $0 < p < 1$*

$$K(\alpha, \beta, p) \langle x, x \rangle \sharp_p \langle x, Tx \rangle \leq \langle x, T^p x \rangle \leq \langle x, x \rangle \sharp_p \langle x, Tx \rangle$$

for every nonsingular element  $x \in \mathcal{X}$ , where  $K(\alpha, \beta, p)$  is defined by (3.3).

#### 4. HÖLDER INEQUALITY

As an application of Theorem 3.1 and Theorem 3.3, we show Hölder type inequalities on a Hilbert C\*-module and its reverses.

**Theorem 4.1.** *Let  $A$  and  $B$  be positive invertible operators in  $\mathcal{L}(\mathcal{X})$  and  $x$  a nonsingular element of  $\mathcal{X}$ , and  $\frac{1}{p} + \frac{1}{q} = 1$ .*

(1) *If  $p > 1$ , then*

$$\langle x, B^q \sharp_{1/p} A^p x \rangle \leq \langle x, B^q x \rangle \sharp_{1/p} \langle x, A^p x \rangle \quad (4.1)$$

or

$$\langle x, A^p \sharp_{1/q} B^q x \rangle \leq \langle x, A^p x \rangle \sharp_{1/q} \langle x, B^q x \rangle. \quad (4.2)$$

(2) *If  $p \leq -1$  or  $\frac{1}{2} \leq p < 1$ , then*

$$\langle x, B^q \natural_{1/p} A^p x \rangle \geq \langle x, B^q x \rangle \natural_{1/p} \langle x, A^p x \rangle \quad (4.3)$$

or

$$\langle x, A^p \natural_{1/q} B^q x \rangle \geq \langle x, A^p x \rangle \natural_{1/q} \langle x, B^q x \rangle. \quad (4.4)$$

*Proof.* Replacing  $x$  and  $T$  by  $B^{\frac{q}{2}}x$  and  $(B^{-\frac{q}{2}}A^pB^{-\frac{q}{2}})^{\frac{1}{p}}$  in (1) of Theorem 3.1 respectively, we have (4.1) of Theorem 4.1. By (4.1) and the symmetric property of  $t$ -geometric mean, we have (4.2). The latter (4.3) and (4.4) are proved similarly.  $\square$

By Theorem 3.5, we have the following weighted version of Cauchy type inequality on a Hilbert C\*-module.

**Theorem 4.2.** *Let  $A$  and  $B$  be positive invertible operators in  $\mathcal{L}(\mathcal{X})$  such that  $\alpha I \leq A, B \leq \beta I$  for some scalars  $0 < \alpha < \beta$ . Then for  $0 < p < 1$*

$$K\left(\frac{\alpha^2}{\beta^2}, \frac{\beta^2}{\alpha^2}, p\right) \langle x, B^2 x \rangle \sharp_p \langle x, A^2 x \rangle \leq \langle x, A^2 \sharp_p B^2 x \rangle \leq \langle x, B^2 x \rangle \sharp_p \langle x, A^2 x \rangle$$

for every nonsingular element  $x \in \mathcal{X}$ .

*Proof.* Replace  $x$  and  $T$  by  $Bx$  and  $B^{-1}A^2B^{-1}$  in Theorem 3.5 respectively. Since  $\frac{\alpha^2}{\beta^2}I \leq B^{-1}A^2B^{-1} \leq \frac{\beta^2}{\alpha^2}$ , the theorem follows.  $\square$

If we put  $p = 1/2$  in Theorem 4.2, then we have the following Pólya-Szegő type inequality on a Hilbert C\*-module which is regarded as a reverse of Cauchy type inequality, also see [8, Theorem 3.3].

**Corollary 4.3.** *Let  $A$  and  $B$  be positive invertible operators in  $\mathcal{L}(\mathcal{X})$  such that  $\alpha I \leq A, B \leq \beta I$  for some scalars  $0 < \alpha < \beta$ . Then*

$$\langle x, Ax \rangle \sharp \langle x, Bx \rangle \leq \frac{\alpha + \beta}{2\sqrt{\alpha\beta}} \langle x, A \sharp Bx \rangle$$

for every nonsingular element  $x \in \mathcal{X}$ .

Next, we show a complementary version of Theorem 4.1.

**Theorem 4.4.** *Let  $A$  and  $B$  be positive invertible operators in  $\mathcal{L}(\mathcal{X})$  such that  $\alpha I \leq A, B \leq \beta I$  for some scalars  $0 < \alpha < \beta$ , and  $x$  a nonsingular element of  $\mathcal{X}$  and  $\frac{1}{p} + \frac{1}{q} = 1$ .*

(1) *If  $p > 1$ , then*

$$\langle x, B^q x \rangle \sharp_{1/p} \langle x, A^p x \rangle \leq K \left( \frac{\alpha}{\beta^{q-1}}, \frac{\beta}{\alpha^{q-1}}, p \right)^{\frac{1}{p}} \langle x, B^q \sharp_{1/p} A^p x \rangle.$$

(2) *If  $p \leq -1$  or  $1/2 \leq p < 1$ , then*

$$\langle x, B^q x \rangle \natural_{1/p} \langle x, A^p x \rangle \geq K \left( \frac{\alpha^p}{\beta^q}, \frac{\beta^p}{\alpha^q}, \frac{1}{p} \right)^{-1} \langle x, B^q \natural_{1/p} A^p x \rangle.$$

*Proof.* Replace  $x$  and  $T$  by  $B^{\frac{q}{2}}x$  and  $(B^{-\frac{q}{2}}A^pB^{-\frac{q}{2}})^{\frac{1}{p}}$  in (1) of Theorem 3.3 respectively. Since  $\alpha/\beta^{q-1}I \leq (B^{-\frac{q}{2}}A^pB^{-\frac{q}{2}})^{\frac{1}{p}} \leq \beta/\alpha^{q-1}I$ , we have (1) of Theorem 4.4. The latter (2) are proved similarly.  $\square$

Next, we discuss Hölder type inequalities in a complementary interval of Theorem 4.1.

**Corollary 4.5.** *Let  $A$  and  $B$  be positive invertible operators in  $\mathcal{L}(\mathcal{X})$  such that  $\alpha I \leq A, B \leq \beta I$  for some scalars  $0 < \alpha < \beta$ , and  $x$  a nonsingular element of  $\mathcal{X}$  and  $\frac{1}{p} + \frac{1}{q} = 1$ . If  $-1 < p < 0$  or  $0 < p < \frac{1}{2}$ , then*

$$\begin{aligned} K \left( \frac{\alpha^p}{\beta^q}, \frac{\beta^p}{\alpha^q}, \frac{1}{p} \right)^{-1} \langle x, B^q \natural_{1/p} A^p x \rangle &\leq \langle x, B^q x \rangle \natural_{1/p} \langle x, A^p x \rangle \\ &\leq K \left( \frac{\alpha^p}{\beta^q}, \frac{\beta^p}{\alpha^q}, \frac{1}{p} \right) \langle x, B^q \natural_{1/p} A^p x \rangle. \end{aligned}$$

*Proof.* Replacing  $x$  and  $T$  by  $B^{\frac{q}{2}}x$  and  $(B^{-\frac{q}{2}}A^pB^{-\frac{q}{2}})^{\frac{1}{p}}$  in Corollary 3.4 respectively, we have this theorem.  $\square$

## 5. WEIGHTED CAUCHY-SCHWARZ INEQUALITY

In this section, we discuss weighted Cauchy-Schwarz inequality on a Hilbert C\*-module. We cite [9] for the case of the Hilbert space operator.

For  $T \in \mathcal{L}(\mathcal{X})$ , we denote the range of  $T$  and the kernel of  $T$  by  $R(T)$  and  $N(T)$ , respectively. A closed submodule  $\mathcal{M}$  of  $\mathcal{X}$  is said to be *complemented* if  $\mathcal{X} = \mathcal{M} \oplus \mathcal{M}^\perp$ . Suppose that the closures of the ranges of  $T$  and  $T^*$  are both

complemented. Then it follows from [13, page 30] that  $T$  has a polar decomposition  $T = U|T|$  with a partial isometry  $U \in \mathcal{L}(\mathcal{X})$  and  $N(U) = N(|T|)$ . Also, we showed in [8, Lemma 6.1] that

$$|T^*|^q = U|T|^qU^* \quad \text{for any positive number } q. \quad (5.1)$$

As a generalization of Theorem 2.3, we have the following inequality.

**Theorem 5.1** (Weighted Cauchy-Schwarz Inequality). *Let  $T$  be an operator in  $\mathcal{L}(\mathcal{X})$  such that the closures of the ranges of  $T$  and  $T^*$  are both complemented. If  $x, y \in \mathcal{X}$  such that  $\langle Tx, y \rangle$  has a polar decomposition  $\langle Tx, y \rangle = u|\langle Tx, y \rangle|$  with a partial isometry  $u \in \mathcal{A}$ , then the following inequality holds*

$$|\langle Tx, y \rangle| \leq u^* \langle x, |T|^{2\alpha} x \rangle u \sharp \langle y, |T^*|^{2\beta} y \rangle \quad (5.2)$$

for any  $\alpha, \beta \in [0, 1]$  and  $\alpha + \beta = 1$ . In particular,

$$|\langle Tx, y \rangle| \leq u^* \langle x, |T|^2 x \rangle u \sharp \langle y, UU^* y \rangle$$

and

$$|\langle Tx, y \rangle| \leq u^* \langle x, U^* U x \rangle u \sharp \langle y, |T^*|^2 y \rangle.$$

Moreover, under the assumption that  $\langle y, |T^*|^{2\beta} y \rangle$  is invertible for  $\beta \in [0, 1]$ , the equality in (5.2) holds if and only if  $Txu = |T^*|^{2\beta} yb$  for some  $b \in \mathcal{A}$ .

*Proof.* In the case of  $\alpha = 0$  or 1, it follows from Theorem 2.1 that

$$|\langle Tx, y \rangle| = |\langle |T|x, U^* y \rangle| \leq u^* \langle x, |T|^2 x \rangle u \sharp \langle y, UU^* y \rangle$$

and

$$\begin{aligned} |\langle Tx, y \rangle| &= |\langle x, |T|U^* y \rangle| = |\langle x, U^* U |T|U^* y \rangle| = |\langle Ux, |T^*|y \rangle| \\ &\leq u^* \langle Ux, Ux \rangle u \sharp \langle |T^*|y, |T^*|y \rangle = u^* \langle x, U^* U x \rangle u \sharp \langle y, |T^*|^2 y \rangle \end{aligned}$$

by (5.1).

In the case of  $0 < \alpha < 1$ , we have

$$\begin{aligned} |\langle Tx, y \rangle| &= |\langle U|T|x, y \rangle| = |\langle |T|^\alpha x, |T|^\beta U^* y \rangle| \quad \text{by } \alpha + \beta = 1 \\ &\leq u^* \langle x, |T|^{2\alpha} x \rangle u \sharp \langle y, U|T|^{2\beta} U^* y \rangle \quad \text{by Theorem 2.1} \\ &= u^* \langle x, |T|^{2\alpha} x \rangle u \sharp \langle y, |T^*|^{2\beta} y \rangle. \quad \text{by (5.1).} \end{aligned}$$

Next, we consider the equality conditions in (5.2). Since  $\langle Tx, y \rangle = \langle |T|^\alpha x, |T|^\beta U^* y \rangle$  and  $\langle y, |T^*|^{2\beta} y \rangle$  is invertible for  $\beta \in [0, 1]$ , it follows from Theorem 2.1 that the equality in (5.2) holds if and only if  $|T|^\alpha x u = |T|^\beta U^* y b$  for some  $b \in \mathcal{A}$ . Since  $|T|x = 0$  if and only if  $|T|^{1/2}x = 0$ , it follows that  $N(|T|) = N(|T|^q)$  for any positive real numbers  $q > 0$ . If  $|T|^\beta (|T|^\alpha x u - |T|^\beta U^* y b) = 0$ , then  $|T|^q (|T|^\alpha x u - |T|^\beta U^* y b) = |T|^{\alpha+q} x u - |T|^{\beta+q} U^* y b = 0$  for any  $q > 0$  and this implies  $|T|^\alpha x u - |T|^\beta U^* y b = 0$ . Therefore we have the following implications:

$$\begin{aligned} |T|^\alpha x u = |T|^\beta U^* y b &\iff |T|^{\alpha+\beta} x u = |T|^{2\beta} U^* y b \iff U|T|x u = U|T|^{2\beta} U^* y b \\ &\iff Txu = |T^*|^{2\beta} y b \quad \text{by (5.1).} \end{aligned}$$

□

If we put  $\alpha = \beta = \frac{1}{2}$  in Theorem 5.1, then we have the following inequality.



**Theorem 5.2.** *Let  $T$  be an operator in  $\mathcal{L}(\mathcal{X})$  such that the closures of the ranges of  $T$  and  $T^*$  are both complemented. If  $x, y \in \mathcal{X}$  such that  $\langle Tx, y \rangle$  has a polar decomposition  $\langle Tx, y \rangle = u|\langle Tx, y \rangle|$  with a partial isometry  $u \in \mathcal{A}$ , then*

$$|\langle Tx, y \rangle| \leq u^* \langle x, |T|x \rangle u \# \langle y, |T^*|y \rangle. \quad (5.3)$$

*Moreover, under the assumption that  $\langle y, |T^*|y \rangle$  is invertible, the equality in (5.3) holds if and only if  $Txu = |T^*|yb$  for some  $b \in \mathcal{A}$ .*

**Acknowledgement.** The authors would like to express their cordial thanks to the referee for his/her valuable suggestions.

## REFERENCES

1. T. Ando, *Concavity of certain maps on positive matrices and applications to Hadamard products*, Linear Algebra Appl. **26** (1979), 203-241.
2. T. Ando and F. Hiai, *Hölder type inequalities for matrices*, Math. Ineq. Appl. **1** (1988), 1-30.
3. J.-C. Bourin, E.-Y. Lee, M. Fujii and Y. Seo, *A matrix reverse Hölder inequality*, Linear Algebra Appl., **431** (2009), 2154–2159.
4. M.D.Choi, *A Schwarz inequality for positive linear maps on  $C^*$ -algebras*, Ill. J. Math. **18** (1974), 565–574 .
5. C.Davis, *A Schwartz inequality for convex operator functions*, Proc. Amer. Math. Soc. **8** (1957), 42–44.
6. J.I. Fujii, M. Fujii, M.S. Moslehian and Y. Seo, *Cauchy–Schwarz inequality in semi-inner product  $C^*$ -modules via polar decomposition*, J. Math. Anal. Appl., **394** (2012), 835-840.
7. M. Fujii, S. Izumino, R. Nakamoto and Y. Seo, *Operator inequalities related to Cauchy-Schwarz and Hölder-McCarthy inequalities*, Nihonkai Math. J., **8** (1997), 117–122.
8. J.I. Fujii, M. Fujii and Y. Seo, *Operator inequalities on Hilbert  $C^*$ -modules via the Cauchy-Schwarz inequality*, to appear in Math. Inequal. Appl.
9. T. Furuta, *Invitation to Linear Operators*, Taylor&Francis, London, 2001.
10. T. Furuta, J. Mičić Hot, J. Pečarić and Y. Seo, *Mond-Pečarić Method in Operator Inequalities*, Monographs in Inequalities 1, Element, Zagreb, 2005.
11. S. Izumino and M. Tominaga, *Estimations in Hölder type inequalities*, Math. Inequal. Appl., **4** (2001), 163–187.
12. F. Kubo and T. Ando, *Means of positive linear operators*, Math. Ann., **246** (1980), 205–224.
13. E.C. Lance, *Hilbert  $C^*$ -Modules*, London Math. Soc. Lecture Note Series 210, Cambridge Univ. Press, 1995.
14. V.M. Manuilov and E.V. Troitsky, *Hilbert  $C^*$ -Modules*, Translations of Mathematical Monographs, 226, American Mathematical Society, Providence RI, 2005.
15. B. Mond and O. Shisha, *Difference and ratio inequalities in Hilbert space*, "Inequalities II", (O.Shisha, ed.), Academic Press, New York, 1970, 241–249.
16. W.L. Paschke, *Inner product modules over  $B^*$ -algebras*, Trans. Amer. Math. Soc., **182** (1973), 443–468.

<sup>1</sup> DEPARTMENT OF MATHEMATICS EDUCATION, OSAKA KYOIKU UNIVERSITY, 4-698-1 ASAHIGAOKA, KASHIWARA, OSAKA 582-8582 JAPAN.

*E-mail address:* [yukis@cc.osaka-kyoiku.ac.jp](mailto:yukis@cc.osaka-kyoiku.ac.jp)